

# Fertility Guidelines for Hops Grown in Ontario (2020 update):

There are currently no Ontario fertility recommendations for hops. The following tables for lime, nitrogen (N), phosphorus (P) and potassium (K) applications are guidelines only and are provided for reference based on similar common crops and published research on hops in other jurisdictions (note: lime, phosphorus and potassium should always be applied according to a soil test).



## LIME:

Studies suggest that the optimal soil pH range for hops is 6.5-7.5. However, as with most horticultural crops, a pH range of 6.0 and above is typically suitable for plant growth. Ground limestone is traditionally used to correct soil acidity and is easiest to apply prior to crop establishment. In the case of perennial crops it can be difficult to incorporate large amounts of limestone after crop establishment. Applying and incorporating split applications of total recommended lime over multiple years will help distribute the lime more uniformly over a pre-existing yard and make it easier when working around perennial crops such as hops. Soil pH and buffer pH will be provided on your soil test.

Note: Full effects of lime may not be realized for up to three years after application.

Table 1: Lime Requirements to Correct Soil Acidity						
(based on OMAFRA Pub 611, OMAFRA Soil Fertility Handbook)						
	Ground Limestone Required (t/ha)*					
Buffer pH	Target soil pH		Target soil pH			
	= 7.0 <sup>1</sup>	= 6.5 <sup>2</sup>	$= 6.0^3$	= 5.5 <sup>4</sup>		
7.0	2	2	1	1		
6.9	3	2	1	1		
6.8	3	2	1	1		
6.7	4	2	2	1		
6.6	5	3	2	1		
6.5	6	3	2	1		
6.4	7	4	3	2		
6.3	8	5	3	2		
6.2	10	6	4	2		
6.1	11	7	5	2		
6.0	13	9	6	3		
5.9	14	10	7	4		
5.8	16	12	8	4		
5.7	18	13	9	5		
5.6	20	15	11	6		
5.5	20	17	12	8		
5.4	20	19	14	9		
5.3	20	20	15	10		
5.2	20	20	17	11		
5.1	20	20	19	13		
5.0	20	20	20	15		
4.9	20	20	20	16		
4.8	20	20	20	18		
4.7	20	20	20	20		
4.6	20	20	20	20		

<sup>\*</sup>Based on Agricultural Index of 75.

Liming to pH 7.0 is recommended only for club-root control on cole crops.

<sup>&</sup>lt;sup>2</sup> Add lime if soil pH is below 6.1.

Add lime if soil pH is below 5.6.

Add lime if soil pH is below 5.1.

#### **NITROGEN:**

Table 2: Nitrogen Guidelines for Hops				
Plant Age	kg N/ha	Notes		
	(lb/ac)			
New Yard	65-135 (55-120)	Apply ~20 kg N/ha (18 lb N/ac) at sprouting (mid-late April depending on growing zone).		
Established Yard	165-225 (150-200)	Starting mid-May, apply no more than 25 kg actual N/ha (20-22 lbs N/ac) in any one application every two weeks until July 1.  Gradually decrease N application rate at the beginning of July through to harvest (5-10 kg N/ha (4-10 lbs N/ac) in July; 2-5 kg N/ha (2-5 lbs N/ac) in Aug).		

#### PHOSPHORUS AND POTASSIUM:

Table 3: Phosphate Guidelines for Hops Based on OMAFRA Accredited Soil Tests for Similar Vegetative Perennials (based on OMAFRA Publication 360, Guide to Fruit Production, 2016-2017)

Soil Phosphorus	Rating <sup>1</sup>	Phosphate (P <sub>2</sub> O <sub>5</sub> ) <sup>2</sup> required (kg/ha)	
(ppm)*		New Plantings	Established Plantings
0-3	HR	140	100
4-5		130	90
6-7		120	80
8-9		110	70
10-12	MR	100	70
13-15		90	60
16-20	LR	70	50
21-25		60	40
26-30		50	30
31-40	RR	40	20
41-50		0	0
51-60		0	0
61-80	ND	0	0
80+	NR	0	0

<sup>\*0.5</sup>M sodium bicarbonate extract test method.

Table 4: Potassium Guidelines for Hops Based on OMAFRA Accredited Soil Tests for Similar Vegetative Perennials (based on OMAFRA Publication 360, Guide to Fruit Production, 2016-2017)

Potassium Soil Test (ppm)*	Rating	Potassium (K <sub>2</sub> O) <sup>2</sup> required (kg/ha)
0-15		130
16-30	up	120
31-45	HR	110
46-60		100
61-80	MD	90
81-100	MR	80
101-120	LR	70
121-150		60
151-180	20	40
181-210	RR	0
211-250		0
251+	NR	0

<sup>\*1</sup> M ammonium acetate extract test method.

### Additional Soil Fertility Resources:

OMAFRA Pub 611, Soil Fertility Handbook: <a href="https://www.omafra.gov.on.ca/english/crops/pub611/pub611.pdf">www.omafra.gov.on.ca/english/crops/pub611/pub611.pdf</a>

Michigan State University Extension, Graph of Weekly & Cumulative Nitrogen Application for Hops in Michigan: <a href="https://www.canrmsu.edu/contentAsset/image/ecdaf5f5-aeff-4dae-8ab5-d89075ec8e00/fileAsset/filter/Resize\_Ipeg/resize\_w/720/ipeg\_q/80">https://www.canrmsu.edu/contentAsset/image/ecdaf5f5-aeff-4dae-8ab5-d89075ec8e00/fileAsset/filter/Resize\_Ipeg/resize\_w/720/ipeg\_q/80</a>

<sup>&</sup>lt;sup>1</sup>HR, MR, LR, RR, and NR denote, respectively, high, medium, low, rare and no probabilities of profitable crop response to applied nutrient.

<sup>&</sup>lt;sup>2</sup> Where manure is applied, reduce fertilizer applications according to the amount and quality of manure.

<sup>&</sup>lt;sup>1</sup>HR, MR, LR, RR, and NR denote, respectively, high, medium, low, rare and no probabilities of profitable crop response to applied nutrient.

<sup>&</sup>lt;sup>2</sup> Where manure is applied, reduce fertilizer applications according to the amount and quality of manure.